



TOGETHER
for a sustainable future

OCCASION

This publication has been made available to the public on the occasion of the 50th anniversary of the United Nations Industrial Development Organisation.



TOGETHER
for a sustainable future

DISCLAIMER

This document has been produced without formal United Nations editing. The designations employed and the presentation of the material in this document do not imply the expression of any opinion whatsoever on the part of the Secretariat of the United Nations Industrial Development Organization (UNIDO) concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries, or its economic system or degree of development. Designations such as “developed”, “industrialized” and “developing” are intended for statistical convenience and do not necessarily express a judgment about the stage reached by a particular country or area in the development process. Mention of firm names or commercial products does not constitute an endorsement by UNIDO.

FAIR USE POLICY

Any part of this publication may be quoted and referenced for educational and research purposes without additional permission from UNIDO. However, those who make use of quoting and referencing this publication are requested to follow the Fair Use Policy of giving due credit to UNIDO.

CONTACT

Please contact publications@unido.org for further information concerning UNIDO publications.

For more information about UNIDO, please visit us at www.unido.org



BAMBOO CULTIVATION MANUAL

GUIDELINES FOR CULTIVATING ETHIOPIAN HIGHLAND BAMBOO

EASTERN AFRICA BAMBOO PROJECT

funded by



executed by



supervised by



ETHIOPIA

Ministry of Agriculture and Rural Development

Federal Micro and Small Enterprises
Development Agency

KENYA

Kenya Forestry Research Institute

BAMBOO CULTIVATION MANUAL

GUIDELINES FOR CULTIVATING ETHIOPIAN HIGHLAND BAMBOO

This project is being financed by the Common Fund for Commodities, an intergovernmental financial institution established within the framework of the United Nations, headquartered in Amsterdam, the Netherlands.

EASTERN AFRICA BAMBOO PROJECT

funded by



executed by



supervised by




ETHIOPIA

Ministry of Agriculture and Rural Development

Federal Micro and Small Enterprises
Development Agency

KENYA

Kenya Forestry Research Institute



Copyright ©2009 by the United Nations Industrial Development Organization.


This manual is based on the work of Dr. Victor Brias and Mr. Tesfaye Hunde, UNIDO Consultants, in cooperation with the project team and under the supervision of UNIDO Project Manager: Mr. Juergen Hierold.

Designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the Secretariat of the United Nations Industrial Development Organization (UNIDO) concerning the legal status of any country, territory, city, or area, or of its authorities or concerning the delimitation of its frontiers and boundaries. The opinions, figures and estimates set forth are the responsibility of the authors and should not necessarily be considered as reflecting the views or carrying the endorsement of UNIDO. The mention of firm names or commercial products does not imply endorsement by UNIDO.


This document has not been formally edited.

CONTENTS

Introduction.....	1
General Remarks About Bamboo.....	4
Ethiopian Highland Bamboo.....	7
Distribution.....	11
Flowering and Seeds.....	13
Propagation.....	14
Sexual Propagation.....	14
Asexual Propagation.....	14
Propagation by means of Offsets.....	15
Setting up a Bamboo Nursery.....	17
Selection of a Nursery Site.....	17
Nursery Size and Layout.....	18
Temporary and Permanent Nurseries.....	19
Nursery Infrastructure and Equipment.....	20
Preparation of the Open Nursery Field.....	21
Greenhouse Structures and Design.....	22
Covered Work Space.....	23
Water Supply and Irrigation System.....	23




Storage Room.....	24
Potting Soil.....	24
Fences.....	25
Nursery Operations.....	26
Potting Containers.....	26
Preparation of Potting Soil.....	27
Seed Propagation.....	27
Cultivation of Wildlings.....	30
Potting.....	31
Fertilizer Applications.....	33
Macro-Proliferation.....	33
Irrigation.....	35
Weeding.....	35
Hardening.....	36
Quality Control At the Nursery.....	37
General Maintenance.....	37
Quality of the Soil Mix.....	38
Quality Control During Repotting.....	38
Phytosanitary Treatments.....	39



Rodent Control.....	40
Planting Bamboo in the Field	41
Selection and Transport of Planting Material	41
Planting Procedure for Nursery Plants.....	42
Direct planting of offset cuttings	43
Replanting.....	44
Intercropping	44
Plantation Maintenance.....	45
Soil Maintenance	45
Weeding.....	45
Pest Control	45
Mulching	46
Clump Management	46
Reducing the Risk of Forest Fires	47
Harvesting and Handling	48
Post-Harvest Treatment	51
Drying and Seasoning.....	51
Mold prevention	52
Preservation.....	53

Non-chemical methods.....	53
Chemical Preservation Methods.....	54
References.....	55
Appendix 1: Data to be recorded Based on Phonological Observations of bamboo.....	57
Appendix 2: Data to be gathered During Vegetative Phase of highland bamboo.....	58



INTRODUCTION

This manual provides basic information about Ethiopia's highland bamboo species, and thereby aims to increase public awareness about this important natural resource. Practical guidelines for cultivating and managing highland bamboo are offered herein to help growers establish new plantations and manage existing bamboo stands.

The highland bamboo species is botanically known as *Yushania alpina*. This species grows naturally in ecological zones of the country between 2200 – 3500 meters above sea level. The coverage of this species in Ethiopia was roughly estimated in 1997 to be about 130,000 hectares. Over the last 10 years, the resource base has been significantly reduced because large areas of indigenous bamboo forests have been cleared for conversion to agriculture. This is the case in the highland areas of the country within a 200 km perimeter of Addis Ababa.

Alpine bamboo has traditionally been used as a material for making fences and water pipes, as well as a variety of handicrafts. The culms are used by cottage industries for woven and plaited products such as basketry, mats, and other decorative items.

In recent years, *Y. alpina* has been used for manufacturing industrial products such as parquet flooring, window blinds, and curtains. The plant is also useful for bioenergy.




A stand of *Yushania alpina* at Injibara (Amhara Region)

Most of the bamboo raw material used in Ethiopia is extracted from natural stands. Although bamboo is a renewable resource, the capacity of this plant to regenerate is hindered by indiscriminate and destructive harvesting methods. The uncontrolled extraction of natural bamboo stands has resulted in reduced productivity and yields as well as deterioration in quality. The resource base has been diminishing over the years.

Plantations of highland bamboo are needed to meet the increasing demand for raw material by industries, especially near Addis Ababa. There is a shortage of bamboo raw materials in some urban cities such as in Addis Ababa, Awassa and Bahir Dar. Raw materials from new plantations can ensure the growth of the industrial sector while providing income to farmers.

Bamboo farming can be a lucrative business. The current price of bamboo poles of *Y. alpina* ranges from 7-12 Birr per piece. Given the growth of the industrial bamboo sector in the country, farmers who take up the challenge of bamboo cultivation can virtually be assured of an extra source of income.

Hopefully, this manual will encourage local farmers to consider the cultivation of bamboo as an option, which will be beneficial not only to themselves but also to industry and the environment.



GENERAL REMARKS ABOUT BAMBOO

Technically, bamboo is a grass belonging to the subfamily *Bambusoideae*. Over 1,200 different species grow worldwide. Various species can reach heights of 30 m and more. About 18 million ha of bamboo are distributed in world forest ecosystems in Asia, Africa, and America.

Unlike most timber, bamboo is a self-regenerating natural resource; new shoots that appear annually ensure future raw material after mature culms are harvested.

Bamboo provides considerable environmental benefits. In many countries, it is used for ecological purposes such as soil stabilization and erosion prevention on hill slopes and verges. It is a very important forestry plant which is harvested from existing natural forests, plantations, and mixed agroforestry systems. Bamboo silviculture is an option for conserving and protecting tropical forests while creating enduring supplies for the wood and cellulose industries.

Bamboo is a multipurpose plant with a myriad of applications ranging from construction materials, furniture, fences, handicrafts, pulp and paper, edible shoots, and animal fodder. In developing countries, it is a basic raw material with numerous traditional uses. It is highly suitable for handicrafts; it can be woven into numerous products including mats, baskets, trays, hats, lampshades, caps, lanterns, etc. Many bamboo products are functional while others serve decorative purposes.

Apart from its manifold uses in cottage industries, bamboo is also widely used in modern wood and paper industries. Governments, research institutions, and private enterprises around the world are taking increased interest in the environmental and economic possibilities of bamboo. In the last decade, there has been a boom of manufacturing industries utilizing bamboo worldwide.

Bamboo is also a source of food. The cone-shaped sprouts that emerge from the ground to form tall poles are edible vegetables when harvested

young. Bamboo shoots generally appear during spring or the early rainy season. When harvested young, they are a crunchy and nutritious vegetable. Young shoots contain up to 90% water and are rich in vitamins, cellulose, and amino acids. They have a high nutritional value, are low in fat and high in fiber content. Young shoots vary in size and weight according to species; the edible content of a newly harvested shoot is usually 30% of its weight. Bamboo shoots are sold fresh but are also canned in brine. They are exported worldwide and constitute a multi-million dollar trade commodity.

For most products, bamboo processing does not require high capital investments but is labor intensive and contributes significantly to employment. Skilled labor as well as attractive designs and fine finishing are very important in making bamboo products for commercial purposes. The utilization of bamboo fences is widespread in tropical Africa. Applications of bamboo for structural construction, walls, ceilings, room partitions, windows, furniture, ladders, etc. that are common in Asia could also be developed in Ethiopia and neighboring countries.

There has been a growing awareness in recent years that bamboo is a vital component of development and an effective means to improve the livelihoods of rural poor people. Over 600 million people around the world generate income from bamboo. Hundreds of millions of people in the world live in bamboo houses. Women and children, many of whom live below subsistence levels in developing countries, harvest a great part of the bamboo that is used.

Bamboo is a natural vehicle for development because rural people generally have adequate access to it. It can be easily grown and harvested in the perimeter of forest areas or under agroforestry schemes. Bamboo agroforestry requires only a modest capital investment and generates steady income to farmers. In many parts of the tropical world, the rural poor are dependent on bamboo for their shelter and daily domestic uses.



Yushania alpina: Ethiopian Highland Bamboo

ETHIOPIAN HIGHLAND BAMBOO

The botanical name of Ethiopia's highland bamboo is *Yushania alpina* (K. Schum.) Lin. The species was formerly classified under the genus *Arundinaria*, and older literature refers to this bamboo species using the name *Arundinaria alpina*. The species is however commonly referred to as Alpine bamboo. The local name of this species in Affan Oromo is *Lemen*, while in Amharic, it is known as *Kerkeha*.

Y. alpina is a tufted, sympodial bamboo with erect green culms ranging from 2 to 19.5 meters in height. The diameter of the culm ranges from 5 to 12.5 cm.



Highland bamboo stand with emerging culms (with milky white culm sheaths) and older (green) culms

The culm sheaths are have a milky white color and are covered in reddish brown bristly hairs. The culm sheaths have a linear acute blade and fimbriate lateral auricles. Branches and foliage culms appear on the upper nodes of the culm. The leaf-blades are glabrous and narrowly lanceolate, and range from 5-20 cm in length and 6-15 mm width. The leaves have conspicuous cross veins, narrowed to a fine bristle at the apex.



Left: Mr. Tesfaye Hunde examines new culms of *Y. alpina*

Right: Detail of sheaths still intact on a new culm

Like all bamboos, *Y. alpina* is made up of an underground axis and an above-ground axis. The underground axis consists of rhizomes, roots, and buds. The rhizomes collect and store the nutrients that sustain the life of the plant.



A new shoot of Yushania alpina

Buds on the rhizomes develop into shoots that emerge from the ground to form a clump of culms. The rhizomes of *Y. alpina* exhibit a sympodial branching pattern which gives the plant a clump forming habit. The clumping habit is evident especially in plants under cultivation. The species however has elongated rhizome necks, and the culms are usually widely spaced, forming a loose or open clump. In dense natural stands, it becomes difficult to distinguish one clump from another.

The bamboo culm is cylindrical and is divided into sections by diaphragms or *nodes*. The section between two nodes is called an *internode*. The culm of *Y. alpina* is hollow but it typically has a thick wall, which makes it ideal for the production of strong panel products.

New culms of *Y. alpina* grow to their full height within 3 months and thereafter develop branches and leaves on the upper nodes. As the culm matures, it lignifies and becomes harder and stronger. Unlike trees, bamboos do not undergo secondary growth. In other words, the culm does not get thicker each year. Rather, as the rhizome system develops

and matures, new and larger shoots emerge annually until the maximum culm size is reached.



Tall culms of *Yushania alpina*

Culms of *Y. alpina* are fully mature 3 years after they emerge from the soil. As the mature culm ages, it deteriorates and eventually dies and rots. The life of the bamboo clump is however sustained by the rhizome system which annually generates new shoots which grow into culms.

Effective management involves systematic but selective cutting of mature culms. Selective harvesting of the crop ensures a sustainable supply of valuable and useful raw material. The removal of mature culms also maintains the vigor of the plant and allows for the continuous generation of new shoots.

DISTRIBUTION

Y. alpina is found in ecological zones of the country between 2200 – 3500 meters above sea level.

The species also occurs naturally in mountain areas in East African countries such as Kenya, Uganda and Tanzania.

Most of the alpine bamboo in Ethiopia grows naturally in highland forests. The uncontrolled exploitation of this resource has resulted in reduced productivity and yields as well as deterioration in quality. In recent years, it has been overexploited creating a shortage of raw material supplies in some urban cities such as in Addis Ababa, Awassa, Bahir Dar, and other cities and towns.

The resource base of *Y. alpina* in Ethiopia was roughly estimated in 1997 to be about 130,000 hectares. Since 1997, no inventory has been taken for this species.

Increased exploitation during the last 10 years has resulted in a significant reduction in the bamboo forest area in the country.

Site Name	Region	Zone	Remarks
Agaro	Oromiya	Jima	Natural stands
Bore	Oromiya	Gujji	Private plantations
Ambo-Shenen	Oromiya	Western Showa	Private plantations
Bale	Oromiya	Bale	Natural stands
Gera	Oromiya	Jimma	Natural stands
Jibat Mountain	Oromiya	Western Showa	Natural stands
Harena Forest	Oromiya	Bale	Mixed forest with bamboo
Degaga Munesa Shashemene Enterprise	Oromiya	Western Aris	Mixed forest with bamboo
Awi Zone	Amhara	Awi Zone(Injibara)	Private plantations
Hagere selam	SNNPR	Sidama	Private plantations
Ameya	SNNPR	Kefecho Shekecho	Mixed forest with bamboo
Baha-Chapa	SNNPR	Kefecho Shekecho	Natural stands
Bonga	SNNPR		Natural stands
Chincha	SNNPR	North Omo	Private plantations
Gada	SNNPR	Kefecho Shekech	Near Masha area usually combined with Yena
Gecha-Masha	SNNPR	Kefecho Shekecho	Use Masha to define this natural bamboo area.
Indibir	SNNPR	Gurage	Farmer-owned bamboo area extends to Jembero.
Indibir-Jembero	SNNPR	Gurage	Private plantations
Jembero	SNNPR		Private plantations

FLOWERING AND SEEDS

Bamboos, like all grasses in the plant kingdom, are seed plants. Flowering and seeding are necessary for reproduction and new generations. Flowering of some bamboos is cyclic (e.g. every 30 or 100 years), but for most bamboos, flowering is unpredictable. The flowering of bamboos is sometimes followed by the death of the plant, but this is not the case for all species.

In the case of *Y. alpina*, documented information on flowering is scarce. The species has not been observed to flower gregariously and no flowering cycle for this species is known. It is however well known that solitary clumps flower sporadically and shed seed. Sporadic flowering is unpredictable, occurs only on individual clumps, and is not necessarily followed by the death of the plant.

In Ethiopia, it has been observed that sporadic flowering of *Y. alpina* is followed by the death of the flowering clumps. When flowering occurs, the clumps gradually dry up, shed seeds, and die. It may take up to two years before seeds are released from the flowering clump. Seeds germinate about two weeks after they fall on the soil. Seeds are not always fertile; if they are, their viability does not exceed one year.

In general, there is still much to learn about the flowering of bamboo, and particularly about the flowering of the species - *Y. alpina*.

It is however clear that the flowering of the species cannot be predicted with any degree of certainty.

Flowering is a rare occurrence. If bamboos start to flower, it is advisable to monitor the event and to collect seeds or wildlings in order to raise and propagate a new generation of planting material.

PROPAGATION

Currently, there are very few plantations of *Y. alpina* in Ethiopia. Most of the highland bamboos in the country grow naturally in forests.

The uncontrolled exploitation of this resource, however, has resulted in reduced yields and deterioration in quality.

Considering the economic importance of *Y. alpina*, the continuity of bamboo supplies for local and urban uses will depend on the establishment of plantations and on the effective management of existing forests.

SEXUAL PROPAGATION

Sexual propagation involves the production of new bamboo plants through seeds. Seeds of *Y. alpina* are viable for a limited time, seldom for more than 10 months. Because of this poor viability, seeds need to be collected and sown in nursery beds without delay. If wildlings or naturally growing highland bamboo seedlings in the forests are available, these can be collected and used as planting materials.

ASEXUAL PROPAGATION

Bamboos can more easily be propagated using vegetative material such as rhizomes and offsets. Propagules are considered to be successfully established when they form new rhizomes and start to develop new shoots.

PROPAGATION BY MEANS OF OFFSETS

The most effective way of propagating *Y. alpina* is by means of offsetting a culm from a clump. Although this method has a high success rate, it is not the most practical method for large scale propagation, nor is it the best way of obtaining planting material that will be directly planted in the field. This is due to the labor intensiveness of the method and the time and logistical costs involved in obtaining and moving a large number of offsets to the field. The method of offset propagation is however very useful when one desires to raise a few clumps in a homestead or farm. It is also an excellent way of obtaining mother plants that can be further macroproliferated at nursery.

The method of offsetting involves separating culms from clumps and shortening them to the node above the first branches. Since the branches of *Y. alpina* typically appear at upper nodes of the culm, the offsets will be very long and heavy, and not practical for transporting to another location. Some preparatory work is therefore advisable before proceeding with actual offsetting.

- 6 to 12 months before obtaining offsets, make a selection of 1 to 2 years old culms that will be used. Culms of this age can be identified by their green color, and their culm sheaths have fallen off.
- The culms should be selected from a healthy and vigorous bamboo clump in the forest or homestead farm.
- Cut down or coppice the culms just above the 4th or 5th node from the ground.
- Remove all foliage and small branches and use them to mulch the clump. This returns nutrients to the soil and is a practical means of clump management.
- The coppiced culms will generate branches in the lower nodes. These shortened culms can be effectively offsetted after they have

developed some branches and foliage which enable photosynthesis and growth.

- At the onset of the rainy season and just before the emergence of new shoots, offsets can be obtained from bamboo stands as outlined below:
- If the above mentioned preparatory activities are carried out, the selected coppiced culms which have developed some branches and foliage at the lower nodes can be used as offsets.
- Dig out about 60 cm below the ground for a rhizome of a one to two years old culm.
- Once a rhizome is exposed, cut back the aerial culm just above the first branches with leaves but not less than 3rd node above the ground level.
- Cut the rhizome off from the parent clump. Avoid injuring the junction of the culm and rhizome, and the underground dormant buds at the base of the culm.
- Cut the roots and soil at least 10 cm away from the rhizome so that the offset includes rhizomes with roots and soil.
- Replant the offset immediately and flood it with water, and mulch it.
- If planted directly on the ground, make sure that the planting hole is sufficiently deep and large to facilitate the easy development of new shoots.
- If planted in a container, make sure that the container is sufficiently large and deep to allow the plant to grow as it would on the ground. Make sure that the container has holes so that there is good drainage.
- Water the plant frequently so that the soil is always moist.

SETTING UP A BAMBOO NURSERY

Establishing a bamboo nursery is a practical solution for maintaining a regular supply of planting material for plantations and forestry projects. The nursery does not require high investments. Basic equipment for manual operations can be used.

SELECTION OF A NURSERY SITE

Several factors should be considered in selecting a site for a bamboo propagation nursery.

Location and Accessibility: If possible, the nursery should be located near a highway or public road to facilitate operations, communication and transport. Ideally, the nursery site should be as close as possible to plantation areas. This will involve less transport time and costs in delivering plants to the field. Plants will undergo less stress during transport the quicker they can be delivered to the field.

Water Supply: This is a critical aspect of a bamboo propagation nursery. The nursery should be located in an area where there is an abundant and permanent supply of water. If the supply of water during the dry seasons is inadequate, a storage tank should be constructed. The amount of water needed depends upon the nursery size, watering frequency, rainfall and climatic conditions, the species to be raised, the quantity of propagation materials, and the method of watering to be employed.

Topography: Local topography is a crucial factor and ideally, the site should slope gently to about 5° so that rainwater can run off without causing erosion. In general, hilltops and valley bottoms are unsuitable; locations on middle to lower slopes are preferable.

Soil: Nursery production requires fertile and well-drained soil with a medium to light texture.

Sun and Shade: Plants in a nursery need a good balance of sunlight and shade. Sites that are heavily shaded throughout the day should be avoided. Partial shading is desirable in the very dry areas to prevent excessive day temperature. Arid areas with desiccating conditions are not suitable for a bamboo nursery.

NURSERY SIZE AND LAYOUT

The size of the nursery depends on its intended production capacity and on the size and age of plants to be grown. A nursery dedicated to the production of young liners or seedlings which are to be delivered to other nurseries will require less space than a nursery which grows and stocks older and larger plants.

On average, the amount of time bamboo seedlings need to remain in the nursery ranges from 8 to 12 months. In general, at least 5000 m² should be allotted for every 10,000 seedlings or young plants that will be raised annually. If seedlings will be grown at the nursery until they are 2 years old, then the area needed for production should be doubled. Adequate space is needed to maintain a stock of growing plants. A crowded nursery will only result in the production of poor stock quality.

The shape or layout of the nursery should be approximately square to minimize the length of the perimeter. This will not only reduce the cost of fencing the nursery but will also enable faster movement of workers from one point of the nursery to another.

The area needed for paths, roads, irrigation, ditches, and buildings should also be taken into consideration, and represents additional space requirements to the entire nursery unit. Having more land than initially required provides an allowance for future expansion of the nursery production area.

TEMPORARY AND PERMANENT NURSERIES

The type of nursery that is established, including all infrastructure requirements, will depend on the duration of cultivation.

If plants or seedlings only need to be cultivated to provide planting material for one large plantation or for a group of small scale farmers in a given area, then only a temporary nursery is usually sufficient.

A temporary nursery can be set up using very basic and low-cost materials that can often be obtained in villages near the site. Concrete structures and sophisticated and expensive tools are not required; simple shade houses using palm or bamboo roofing are suitable in most cases.



A temporary bamboo nursery in the Ethiopian highlands

However, if the nursery is intended to be used for a period of at least five to ten years, then permanent and durable structures should be built on site. The nursery should also be easily accessible by road at all times of the

year. A permanent nursery is needed when mass propagation of planting stocks will be undertaken year after year.



Nursery of the Forestry Department in Addis Ababa

NURSERY INFRASTRUCTURE AND EQUIPMENT

The infrastructure of a nursery generally includes an office and a meeting room for workers, a warehouse or storeroom, a potting shed, a compost area, lath houses, shade tunnels, and an open-air container ground. The number and size of the structures will depend on the production capacity of the nursery.

Polyester ground cover cloth for the open air ground for plants in container or polybags is used in many professional nurseries since this prevents plants from rooting into the ground. This material is costly however, and is optional. Such an investment is however worthwhile when the nursery is intended for commercial purposes and investment costs will be recuperated through the sale of plants.

Irrigation in the nursery is essential. Irrigation may be done manually, but it is advisable to invest in a sprinkler system to ensure proper irrigation, especially in larger nurseries.

Tools such as spades, picks, pruning scissors, and wheelbarrows are essential equipment for manual operations in every nursery. Such equipment is affordable and does not demand a high investment.

Construction of the nursery can be divided into two phases. The first phase includes planning the site layout, clearing the land, leveling of the nursery grounds, excavating gutters and ditches, laying foundations, preparing pathways, installing piping and/or irrigation hoses, and primary installment of utilities. The second phase covers the construction of the buildings, lath houses, shade tunnels, and sheds.

PREPARATION OF THE OPEN NURSERY FIELD

Clearing and leveling of the land is the first step. The open field should be at a gentle slope of about 5° to ensure water run-off. The ground must be flat to prevent the water from accumulating in puddles. This is imperative because the bamboo plants will be cultivated in polybag tubes, which must never be submerged in water. The drainage system must be well planned. Gutters and trenches should be dug out to prevent water from flooding the container field.

As mentioned, using ground cover cloth is a proven way of reducing the risk of grasses and weeds from infesting the open field production area. The synthetic woven fabric is made of tough UV stabilized polypropylene yarns and is designed to inhibit weed growth and improve the appearance of the nursery. It blocks sunlight and thereby prevents weeds from infesting the potted plants while allowing water and nutrients to freely pass through the soil. The use of ground cover fabric eliminates the need of applying large quantities of herbicides to the soil prior to bringing in the potted plants. While weed control through the application of herbicides implies a lower initial cost, it however involves work and maintenance all year long. The use of a ground cover cloth involves a moderate

investment, but weed control will be more effective and requires less maintenance and labor. This option should therefore be considered depending on the purpose of the nursery and production scale that is forecasted.

GREENHOUSE STRUCTURES AND DESIGN

A greenhouse can be established without costly materials and equipment that are commonly used by nurseries in temperate regions. In tropical countries such as Ethiopia, there is only a slight variation in seasonal and nocturnal temperatures. This eliminates the need for expensive climate control systems. On the other hand, it is essential that the greenhouse structure is designed such that good cross ventilation can be provided by openings in the sides of the greenhouse.

The greenhouse structure can consist of a metal or wooden framework. The aim is to build sturdy structures that provide protection from wind, rain and excessive sunlight while providing very good natural ventilation.

A lath house or open-sided structure can be used to provide shade to young bamboo plants. The lath house protects container grown plants from high temperature and high light intensities. It is effective in reducing moisture stress and reduces irrigation requirements.

The roof of the lath house should be spaced with laths or slats, spaced with gaps as wide as the slats themselves, providing approximately 50% shade. Bamboo or palm thatching may be used as a low-cost roofing material for the lath house. The walls are ideally made of cloth also providing at least 50% shade. The walls of the lath house should be inclined so they can resist wind. Doors can be made by cutting the cloth of the walls and can be kept closed using hooks.

A lath house has many uses. It is, first of all, useful for maintenance and as a shaded area to safeguard plant. It is also useful as an area for propagation and transplanting young plants.

Alternatively, or in combination with lath houses, shade tunnels can be used at the nursery. Shade tunnels reduce light levels for the plants in cultivation protecting them from leaf bleaching or leaf scorch in strong sunlight. At modern nurseries, shade tunnels are usually constructed using steel hoops which are covered with polyethylene film or shade cloth.

A low-cost shade tunnel can however be made using a combination of wood and bamboo strips, in combination with white polyethylene film. Flexible bamboo strips can be joined together and shaped into hoops. The bamboo hoops are then positioned on the ground to form a tunnel framework that is supported by round wood or bamboo poles. The framework is then covered with a polyethylene film to provide shade and protection to the plants. This low-cost alternative will have the same effect as more expensive metal structures. It may not be very durable, but replacing or repairing any damages to the tunnel structure will involve very low costs.

Ventilation in the shade tunnel is achieved by creating openings on the sides and at both ends of the tunnel. Cross ventilation through these openings will enable the structure to withstand strong winds.

COVERED WORK SPACE

It is useful and advisable to provide a sufficiently large covered work space where propagation activities can be carried out. The design of the work space should be simple and functional. Benches made of wood can be used to facilitate manual operations such as dividing plants and potting. Aisles space is necessary to allow easy movement between the benches, while at the same time maximizing the production space.

WATER SUPPLY AND IRRIGATION SYSTEM

Provisions should be made to allow for water harvesting during the rainy season in order to have sufficient water supplies during dry periods. A

large tank can be used for this purpose; alternatively, a water reservoir can be excavated.

It is advisable to have a basic irrigation system including a deep well, pump system, and sprinkler system with controllers. This equipment involves a modest investment, which can go a long way in maintaining healthy and vigorous plants in the nursery.

The alternative to investing in an irrigation system is to have a disciplined workforce that diligently waters the plants in the nursery every day. For a small nursery, manual watering of plants is possible, but for a nursery with a production capacity of thousands of plants, investments in a pump and sprinkler system are necessary to ensure proper irrigation and to minimize plant mortality.

STORAGE ROOM

A small building or shed is needed as a stock room for storing pesticides, fertilizers and tools. The stock room must be spacious enough to stock all materials. It should be well insulated to protect all materials from rain and excessive humidity.

POTTING SOIL

An area should be allocated for stocking components of the soil mixture. There must also be a leveled area to stock organic material for the composting process.

The amount of soil mixture needed will obviously depend on production activities. During periods of intensive potting, several cubic meters of soil mix may be required per day. The soil mixture can be prepared on site using traditional techniques. Making the potting soil involves using the right proportion of materials to create a good balance between water and oxygen allowing for good porosity and drainage. Soil fertility and plant growth can be stimulated using locally available organic inputs such as

cattle and chicken manure, fish residues, groundnut shells, bran, and rock phosphate. In addition to providing nutrients, organic fertilizers also help to conserve water. The soil mixture should have good drainage and a pH value of 5.5, which is perfect for bamboo. Slow release fertilizers, such as Osmocote, can be used to enrich the soil mixture. This fertilizer will supplement the mineral nutrients of the plants in cultivation.

FENCES

A fence should be erected around the perimeter of the nursery to protect the nursery area from domestic and wild animals. The fence should be about 2 meters tall and may incorporate the use of barbed wire for added security.



NURSERY OPERATIONS

Operating a nursery efficiently requires a strong attention to detail and a high level of quality control. Young plants are like infants and need tender loving care so that they grow healthy and strong. The nursery manager or supervisor needs to strictly enforce standard operating procedures to ensure that all the plants in production receive adequate care and attention. Every stage in production, from potting, propagating, weeding, irrigating, fertilizing, etc., is crucial to the survival and vitality of the nursery stock. Staff working at the nursery must diligently perform their duties in the service of the young plants. Nursery workers should be selected carefully based on their ability and dedication to propagate and maintain healthy bamboo plants.

POTTING CONTAINERS

A sufficient quantity of containers or polybags of various sizes should be available at the nursery to fulfill potting requirements. The potting containers should be large enough to allow root development of plants of different sizes.

For seedlings, polythene tubes with 20 cm diameter x 40 cm height x 0.04 mm thickness are sufficient.

Larger polybags with a 40 cm diameter x 50 cm height will be needed to transplant the seedlings once they have grown and have started developing new shoots.

Failure to transplant growing plants to larger containers will impede root development and shoot generation, and inhibit their overall growth.

PREPARATION OF POTTING SOIL

The best organic soils should always be used for cultivating nursery plants. As noted previously, a nursery should be situated near a supply of fertile soil.

A potting mix consisting of 50% forest soils, 25% sand soils and 25% other organic manures is ideal for potting bamboo.

Soil should be sieved to remove large lumps and stones. If necessary, fertilizer should be added in granular form when preparing the soil mixture.

If the soil mix contains many weed seeds, it is desirable to fill containers 3-4 weeks before sowing so that the seeds can germinate and be eliminated in advance.

SEED PROPAGATION

In general, bamboo seeds should not be stored for a prolonged period of time. They have a short-lived viability and their capacity to germinate rapidly begins to deteriorate after two to three months. Because of this poor viability, it is advisable to collect and sow the seeds without delay.

Seeds may be sown in a nursery bed or in polyethylene tubes filled with soil and then covered with a thin layer of soil.

Seeds of *Y. alpina* should be sown either by dibbling with a stick or finger to a depth equal to its shortest dimension, or laid on the surface and covered to this depth.

The planted seeds should be maintained in a shaded area and protected from direct sunlight. This is the favorable environment for high germination. In most cases, not all seeds will germinate, but numerous

seedlings will develop provided that a generous amount of seeds have been planted.

The seeds should be watered daily and must be kept moist at all times. Watering should be done carefully using a fine rose can. Watering is best done in the morning or mid-afternoon, but never at mid-day to avoid the risk of scorching.



A seedling of *Yushania alpina*

Very young bamboo seedlings (as well as small culm cuttings) are very susceptible to sun scorch. Shade capable of blocking up to 75% of direct sunlight should be provided to protect the plants.



Highland bamboo seedlings potted in polybags

Seedlings should be mulched during initial months. Mulching protects them from intense heat, strong winds, and the impact of raindrops. It also helps prevent the rapid evaporation of soil moisture. Dry grass, leaves, and straw are suitable materials for mulching. The kind and amount of mulch will depend on the time of sowing, and rate of growth of seedlings.

Seedlings that have grown to a height of about 5 cm should be gently pricked out of the seedbed and transplanted into polyethylene tubes. After 8-12 months from the date of transplanting, good-sized transplants can be obtained.




Highland bamboo seedlings at a nursery in Amhara region

CULTIVATION OF WIDLINGS

Profuse natural regeneration of *Y. alpina* usually occurs on the forest floor after bamboo flowering. Ripe seeds fall on the ground and germinate on site. Weeds and inter seedlings competition usually affects the establishment of most of these young plants.

Patches of wildlings in a two-leaf stage may sometimes be found on the forest floor after flowering has occurred. In case bamboo seeds are not available, these wildlings can be cultivated and used as propagules at the nursery. The wildlings should be carefully collected by scooping the soil on which they stand and quickly taken to the nursery and transplanted on shaded soil bed. Once the plants have grown to a height of 5cm, they should be individually pricked out and transplanted into polybags.



A bamboo seedling with developed roots

POTTING

Potting of propagules should be done carefully to ensure a high survival rate at the nursery.

Polythene tubes or polybags are the commonly used potting containers. Polybags with a size of 40 cm x 50 cm are suitable for small bamboo plants and generally provide sufficient space for the development of roots, rhizomes, and new shoots.



Highland bamboo seedlings potted in polybags

If the polybags do not have holes at the base, it is necessary to perforate some holes to provide drainage and thereby prevent the roots from being waterlogged.



Potted seedlings properly arranged in the nursery

A light potting mixture with good drainage should be used at all times. Heavy soils should be avoided since they will constrain the growth and development of the young plants.

The potting procedure involves the following steps:

- Fill about 1/3 of the polybags with the soil mixture.
- Place the roots and rhizomes of the plant in the center of the polybag and fill up the remaining portion of the polybag with soil ensuring that the base of the stems are on the level of the soil.
- Compact the soil in the polybag moderately.
- Water the newly-potted plants.
- Place them in a shaded area or in the lath house and ensure they are regularly watered.

FERTILIZER APPLICATIONS

Fertilizer application may be done in the seedbed in conjunction with watering long before transplanting. This is done by dissolving complete NPK fertilizer (14-14-14) or (15-15-15) at the rate of 10g per liter of water.

Fertilizer may also be applied at the time of transplanting. After filling the pot with soil, a pinch (approximately 0.25g) of complete fertilizer is dropped at the hole where the seedling will be transplanted.

If available, slow release fertilizers such as Osmocote can be used for plants that are at least 1 year old.

MACRO-PROLIFERATION

When seeds or wildlings are not available and a large quantity of planting materials is required, the method of offsets should be integrated in a

nursery macro-proliferation program. Offsets taken from the field should be immediately taken to a nursery and planted in large containers with drainage holes. Macro-proliferation involves the method known as clump division.

Division of plants on site provides a means for lowering overall costs and increasing margins. Care must however be taken that this process is planned and supervised in order to maintain a high level of quality.

At the nursery, the following procedure should be followed:

- Offsets placed in containers should be maintained and regular weeding is necessary.
- When the offset has developed at least 2 new culms, it can be divided into at least two plants.
- Division should be done prior to the development of new shoots in the spring.
- Prepare new containers filled with a light potting mixture.
- Remove the plant from its container and shake off the soil so that the rhizome structure can be examined.
- Divide the clump into 2 or more plants following the same method used in offset propagation.
- Replant the divided plants in new pots with a loose potting mixture and water and mulch them regularly.
- One plant should normally yield at least 2 new plants every year.
- Always maintain a stock of plants that are regularly propagated and divided at the nursery.
- To ease delivery of plants to plantations, the plants can be shortened by cutting the stems above the first nodes with leaves and branches.

IRRIGATION

Irrigation is a critical factor for the production of healthy plants. A deep well or water tank must be relied upon as a source of water if the nursery is not located along a lake or river. A pump and sprinkler system will allow for effective watering on a regular basis. Young bamboo plants need water daily; without water the soil humidity in the containers is reduced and the plants can become dehydrated. The plots of growing plants should be controlled on a daily basis. It is also necessary to ensure that the irrigation system is functioning properly so that plants in all areas of the nursery are watered. In the event that the irrigation or sprinkler system fails, plants should be watered manually. Failure to water plants will result in heavy losses.

WEEDING

Weeds compete with bamboo plants by absorbing nutrients, water, and sunlight. Weeds have rhizomes which sprout again if not removed. Measures to control weeds include a thorough preparation of the soil before sowing and the use of sowing media and manure which are free of weed seeds. The removal of weeds growing within the vicinity is also strongly advised in order to lower the occurrence of seed dispersal by wind or water.

The following are recommended practices in weeding:

- Weeding should be done thoroughly, systematically and regularly.
- When removing weeds from soil, no portion of the root system should be left behind.
- Weeding should be done only when the soil is moist.
- Weeds that have been removed should be properly disposed off in a rubbish heap.

- Wherever practical and convenient, mechanical weeding tools may be used.

HARDENING

After being transplanted or potted, the seedlings and cuttings are kept under 50% shade. But as soon as transplants have recovered and commenced growth, they should be exposed to more sunlight and gradually hardened. Hardening is achieved by progressively exposing the plants to more adverse weather conditions similar to those in the planting field. By the time seedlings and cuttings are ready to be planted out in the field, their roots should be well developed and have healthy green foliage, which are prerequisites for shoot development. At least one month before planting out, they should be bared to full sunlight and moved to the open field nursery where they are maintained up to the time they are taken out of the nursery and transported to field planting sites.

Fertilizer should not be applied during the hardening period, especially two months before field planting to prevent the development of succulent tissues.

QUALITY CONTROL AT THE NURSERY

Quality control procedures should be applied from the moment of construction and installation of the nursery up to the time that plants are delivered to customers. The levels of quality control are described below.

GENERAL MAINTENANCE

Throughout the culture, it is necessary to regularly control plants and check on their root and shoot development.

If the plants stay in the nursery longer than anticipated, they must be pruned to a height of 40-50 cm, which can be done using hedge-trimmers.

It is also necessary to make inventories and to make sure that plants are properly labeled and marked. This is crucial especially if the nursery cultivates different bamboo varieties.

The nursery should be kept clean and orderly. Equipment and tools should be regularly inspected, maintained and repaired when necessary. The structures, paths and the plots should be periodically inspected to ensure that they remain in good condition.



Using labels facilitates the identification of species at the nursery.

QUALITY OF THE SOIL MIX

The pH of the soil mix must be controlled and brought to an optimum value. At pH of 5.5, the bamboos assimilate the nutritive elements better and show improved growth. If necessary, the pH of the potting soil mixture can be raised using chemical products such as Calcium carbonate and Calcium sulfate.

QUALITY CONTROL DURING REPOTTING

During repotting, young plants must be protected from direct sunlight and dry winds. This is best achieved by performing the potting activities under a shaded area and immediately transferring the plants to the lath house or shade tunnel.

Careful attention must be paid to the plants during the first three weeks that follow the repotting operation. Initial irrigation must be abundant,

and in the weeks following the repotting operation, irrigation must be frequently done.

PHYTOSANITARY TREATMENTS

Controlling the weeds is a regular maintenance task. If the potting soil components have not been disinfected, there will be too many weed seeds. Effective weed control and reduced manual labor for weeding can be achieved by using anti-germinative herbicides such as Cent 7, which has isoxaben as its active agent.

The first anti-germinative treatment should be carried out just after repotting, at a rate of 6 liters/ha. Three months later, a second treatment must be done, just before the plants are brought to the fields. Nevertheless, a certain amount of manual weeding will be needed during the time the plants are in the nursery. In spite of reducing the time spent in weeding, herbicides cannot completely eliminate weeds and certain weeds manage to break through the Cent 7 film.

The paths between containers in the open field nursery can be treated with a solution of Roundup and Decimax. These herbicides have a systemic effect, penetrate the roots, and have a persistent action in the soil. Herbicidal treatments can be applied 4 to 5 times per year depending on weed infestation.

Bamboos only have a few natural enemies and they normally do not suffer from attacks or diseases when planted in the fields. Nevertheless, a regular control can help to avoid invasions and infections, as well as to determine the tolerance level to any problems. In the nursery, however, certain preventive and curative treatments will be necessary. Following the soil preparation in the nursery and before setting the plots, a treatment against ants and termites must be carried out. Spraying of the soil with Decis (deltamethrine) at 15 g/m² or with Diazinon (240 g/l) can be very effective.

The phytosanitary treatments in the nursery will be essentially of a curative nature, as in the case of the borer beetles that attack the young culms (*Estigmena chinensis*, *Conarthus jansoni*) or leaf rollers (*Pyrausta coclesalis*). These pests occur only occasionally.

Other treatments must have a preventive nature, to avoid the pests commonly found in the nurseries, such as aphids. The use of an insecticide such as the Pirimor will provide a good control over aphid populations. This insecticide has a persisting action and remains active for months; this helps to reduce the number of treatments while protecting the culture.

Treatments with fungicides will be applied when common cryptogamic diseases arise (e.g. *septoriosis* and *fusariosis*). Serious diseases such as the *Rhizoctonia solani* can be avoided by utilizing good potting soil, but frequent controls must nevertheless be carried out.

RODENT CONTROL

Rodents may cause damage by eating the shoots and roots of the bamboo seedlings. Control of these animals can be achieved by laying traps or poisonous baits, or through other local methods.

PLANTING BAMBOO IN THE FIELD

Prior to planting bamboo in the field, a planting scheme which takes into consideration the size and growth habit of the species has to be determined. For *Y. alpina*, the recommended spacing is 7x7m or 204 plants per hectare. This distance provides sufficient space for intercropping and allows greater ease of movement for maintenance and harvesting activities. A 5x5m layout with 400 plants per hectare may however be used for riverbank and gully stabilization, or when intercropping of cash crops is not intended. Field planting can be done either by direct planting of culm offsets or by using nursery-raised plants.

As mentioned earlier, the use of culm offsets for a large plantation is not practical. The preferred type of planting material is bamboo raised at a nursery.

SELECTION AND TRANSPORT OF PLANTING MATERIAL

Plants that are used for planting in the field should be hardened at the nursery before transporting them to the planting site. Plants which are very young and delicate have a lower chance of survival in the field. It is important to select plants that have well developed roots and rhizomes. Such plants will be able to absorb nutrients from the soil and will be able to adapt to the harsher conditions of the field more easily than plants which are still developing roots.

Plants at the nursery which have very long stems may be trimmed down to a height of 50 cm. It is however essential to ensure that the plants have sufficient foliage. Smaller plants with vigorous roots will require less energy for surviving in the field than taller plants.

When transporting plants to the field, they should be handled carefully. The plants should be watered thoroughly prior to transport. They should be loaded and unloaded from the transport vehicle in such a way that no damage is caused to them.

Upon arrival at the field site, the plants should be watered regularly up to the time that they are planted.

PLANTING PROCEDURE FOR NURSERY PLANTS

The following guidelines are suggested to aid the establishment of a productive plantation.

- In selecting the plantation site, check the quality of the soil. Bamboo can grow well on most soil, but deep porous fertile soil with high moisture content and a pH of 5.5 is preferable.
- Good soil drainage is very important. Verify that the land is not prone to flooding. Bamboo does not perform well on waterlogged soils. It is therefore preferable for the plantation to be situated on moderate slopes.
- Clear the land of all weeds and unwanted vegetation. Burning may be necessary during the dry season.
- Carefully plan the layout of plantation so that the planting holes are placed at the specified distances and intervals.
- Plan the activities so that the plantation layout is completed at least 2 weeks before planting.
- The planting holes should be positioned in a north-south orientation. This will provide an optimal distribution of sunlight to all the plants.
- Planting holes with a diameter of 1m and a depth of 60cm should be dug and evenly spaced out according to the plant spacing of 7m x 7m.
- Planting should coincide with the start of the rainy season. If available, organic fertilizer or manure should be placed into each hole and mixed with topsoil. The plants should be planted vertically in an erect position and the hole should be properly covered and mulched.



A large planting hole facilitates the growth of roots and rhizomes.

DIRECT PLANTING OF OFFSET CUTTINGS

Direct planting of offsets of *Y. alpina* in the plantation may be done in small plots or homestead farms. Planting should be conducted at the beginning of the rainy season. The selection and preparation of offset cuttings for direct planting follows the same procedure as that of planting stock for nursery-raised cuttings except that cuttings are directly planted in the field pits without potting.

The procedure for direct planting is as follows:

- Haul the offset cuttings to the planting site.
- Loosen the soil in previously prepared planting pits/holes.
- Place the cuttings in the hole in a vertical position. The lowest node of the culm offset should be above the ground.

- Position the cutting at the center of the planting hole and fill up the pit with soil, ensuring that the culm stands firmly in place.
- Water the soil thoroughly and mulch around the planting hole.

REPLANTING

Not all transplanted seedlings and cuttings will survive the new environment. The plantation should therefore be visited regularly to check on the survival of plants. Dead seedlings and cuttings should be replaced. Replanting should be done immediately at the start of the rainy season.

INTERCROPPING

During the first two years from the time of planting the bamboo, intercropping of cash crops and vegetables may be done between the rows of bamboo. Intercropping serves several purposes. Vegetables provide a source of income to farmers. When cultivated between rows of newly planted bamboo, vegetable crops provide greater stability to the soil and help control erosion. Intercropping also creates an incentive for farmers to control weeds and pests in the plantation. In order to care for their vegetables, farmers will be involved with maintenance activities that are favorable to the growth of the bamboo plants.

When the bamboo canopy has developed, sunlight will be fully absorbed and the cultivation of vegetables will no longer be viable. During the third year after planting the bamboo, leguminous species may be planted to serve as ground cover in between the lines of bamboo.

PLANTATION MAINTENANCE

Maintenance activities during the first 2 years after planting must focus on protecting the young plants from competing vegetation and pests. After the second year, maintenance activities are concentrated on clump management.

SOIL MAINTENANCE

During the first year, it is advisable to loosen the soil around the plant to improve soil aeration. Doing this about twice a year for each plant will enhance growth. Care should however be taken not to disturb the rhizome system of the plant.

WEEDING

The growth of bamboo plants can be hampered by weeds and competing vegetation. It is very important to control and arrest the growth of weeds around each bamboo clump. Failure to do so will invariably result in poor root and stem development in the young bamboos. An area within a radius of 60 cm around each plant should be cleared off of all weeds and vegetation.

PEST CONTROL

The presence of pests and grazing animals should be thoroughly controlled. *Y. alpina* is palatable to animals, especially in dry grazing. It may be necessary to carry out protection against goats and antelopes by partial or total fencing of the plantation. Every available means should be taken to prevent animals from grazing in the plantation. In small homesteads, fencing is a solution, but for a large plantation, it is costly.

Careful supervision in this regard is therefore crucial. It is necessary to patrol the plantation, check for damage, seek the cause, and find suitable means to eliminate the problem.

MULCHING

Mulching is a proven way of improving the growth of bamboo. In drier areas, with rainfall less than 1000 mm, mulching around plants greatly encourages growth through reduced evaporation of soil water. Mulching is achieved by uniformly spreading a layer of leaf litter or other organic material on the surface of the soil around the bamboo clump. Mulching is an effective way of preventing weed growth. It helps conserve soil moisture and contributes organic nutrients to the plant.

Mulching is absolutely necessary for the production of good quality bamboo shoots. The mulch protects young shoots from direct sunlight and keeps them moist, thus allowing them to grow to an optimal size without hardening and losing their edible quality.

CLUMP MANAGEMENT

The proper maintenance of the clump not only improves productivity but also eases the job of the plantation worker. Clump management is partly a maintenance task and partly a result of harvesting. As a maintenance activity, it involves removing unwanted culms to prevent clump congestion. This is particularly necessary with densely tufted species such as *Y. alpina*.

About 90% of new culms emerge in the outer borders of the clump. New shoots and culms cause the clump to widen in diameter. Culms at the periphery of the clump are generally new or young, while older culms stand towards the interior of the clump. Understanding this is important for maintaining a bamboo clump such that its productivity and vigor is maximized.

In maintaining a bamboo clump, it is necessary to extract the oldest culms in the interior of the clump. This is facilitated by creating an opening in the clump and shaping the clump in the form of a horseshoe or “C shape”.

Unless properly managed, clumping bamboos tend to get congested resulting in deterioration both in quality and quantity. It is difficult to extract culms from congested clumps. Preventing clump congestion is important so that harvesting can be done with greater ease. Thinning the clump is essential to provide space for the emergence of new shoots. It is sometimes necessary to sacrifice a few young culms in order to allow for better shoot production in the clump.

Removal of old and rotting culms is also necessary to promote the healthy growth of shoots and new culms. Special attention should be placed on rotting in the stubs of culms that have been harvested. If rotting becomes apparent, it is advisable to dig around the stub and completely remove it. Likewise rotting culms should be extracted. Symptoms of disease or fungal infections should be noted and a plant pathologist should be advised for possible remedies and control measures.

REDUCING THE RISK OF FOREST FIRES

Forest fires are a potential risk to highland bamboo plantations and natural stands, especially during the dry season. Dry leaves, branches and twigs, as well as dead culms may catch and spread fire throughout a large area. Proper mulching of dry leaf litter with humid organic material around the clump aids in preventing the spread of fires.

Firebreaks should be established to safeguard the plantation area. Corridors 10-15 m wide are usually sufficient to stop fire from spreading throughout the plantation.

HARVESTING AND HANDLING

Harvesting of *Y. alpina* should be done selectively according to the age and maturity of the culms. Systematic and selective cutting of mature culms assures the continuous production of young shoots, which is an index of annual yield or increment. The implementation of plantation management strategies can help sustain the regenerative characteristics of bamboo and thereby provide an enduring supply of raw material for industries.

The bamboo plantation will be managed effectively if the exploitation is regulated on a sustainable yield basis. In other words, the clump should never be over harvested or clear cut. New culms as well as 1 to 2 years old culms should not be harvested. A few 3 years old culms should also be left standing so that the clump remains robust and so that harvesting can be performed annually. Following this method, culms are left standing on the clump until they mature, after which, they may be harvested.

A newly established plantation of *Y. alpina* should normally be ready for first harvesting after 5 to 6 years from the time of planting. Thereafter, cutting of mature culms can be done annually or at predetermined intervals of years, according to the management plan and the end use of culms.

The cutting cycles and methods of extraction of culms from a clump should be implemented as an integral part of the management system of the entire bamboo plantation as well as natural stands. Achieving sustainability in terms of culm production will therefore depend on how effectively culms are selected for harvesting, and on how the extraction from the clump is carried out.

Harvesting is a labor intensive operation and it is necessary to make good arrangements with plantation workers so that harvesting operations are not delayed.

The following rules apply for harvesting culms of *Y. alpina* in particular.

- Harvesting of mature culms may begin 5 to 6 years after planting.

- Harvest culms only during the dry season. The starch content of bamboo is lower during periods of dryness. Lower starch content in the culms will make them less susceptible to attack by borers.
- Harvesting should be selective: only mature culms which are 3 to 4 years old should be harvested.
- In a clump, new culms are normally produced outwards - towards the periphery of the clump, and the older culms stand towards the center. Harvesting of culms therefore should be from the center and not at the periphery of the clump. This makes it necessary to maintain clumps in the shape of a horseshoe, keeping the apex towards the side where the new culms are emerging. The open end of the horseshoe facilitates entry inside the clump for extracting mature culms.
- Plan the cutting operation to avoid harming young culms. New culms that attain an average height within the first few months are soft and may collapse unless supported by mature erect culms. A few older culms should be left in the clump after cutting, seeing to it however that congestion is under control.
- Use very sharp tools. It is highly advisable to disinfect harvesting tools using bleach. This lowers the risk of infecting the plants.
- Do not cut young culms unless congestion in the clump prevents the cutting of mature culms.
- Cut each culm between 15 cm to 30 cm from the ground or just above the first node from the ground level. This is necessary so that water does not accumulate in the protruding internode. The accumulation of water may result in rotting and invites insects to lay their eggs.
- Never clear-cut an entire clump unless it has been verified to be seriously infected by a disease.
- Never harvest culms during the rainy season!

- Mulch each clump after harvesting using branches and leaves of harvested culms. These should be neatly piled around the clump to provide organic material to enrich the soil around the clump.

If the plantation is situated near a river, the culms may be allowed to soak in water for a few weeks to aid in the removal of starch and protect them from beetle attack. Otherwise, they should be hauled to an area where they are sorted and air-dried. Good practices to enable drying will help minimize losses due to bio-degradation of the culms. The large culms should be stacked horizontally on parapets where there is good air circulation. Smaller culms may be piled horizontally at a 60° angle to form a “tepee shape” allowing air to circulate around them to aid the drying process. After drying, the whole culms or culm segments are sold in local markets.



POST-HARVEST TREATMENT

Bamboo culms are vulnerable to decay and attack by fungi or insects, especially powder post beetles. Such attacks reduce the natural durability of bamboo and diminish its value and utility.

Post-harvest treatments can help to reduce the risk of decay and attack by pests and thereby increase the useful life and value of bamboo culms.

Depending on the end use of the culm, several methods of preservation may be applied to culms prior to their sale or industrial utilization.

DRYING AND SEASONING

Drying culms is common in the processing of bamboo culm for most uses. Culms are also subjected to seasoning prior to machining, processing, and finishing products that are durable, stable and of a high quality.

Bamboo culms can be air-dried with or without sunlight or they may be kiln-dried. Air-drying is more common than kiln drying since it is more economical. Bamboo culms may be split into halves to speed up drying operations.

Bamboo culms may be thoroughly air-dried in well-ventilated shade for several weeks. Drying can be done by letting the culms stand in a covered area with good air circulation (as shown below).



A large planting hole facilitates the growth of roots and rhizomes

Drying can also be done by stacking culms horizontally on racks. In drying large quantities, the butts and tops of the culms are placed alternately, and then tied in bundles to prevent bending.

Straightening green culms without application of heat requires several weeks. This is done under the shade, either by suspending the freshly-cut curved culm by the tip and attaching a weight at the other end, or by laying the green culm on a flat surface and applying sufficient pressure over the culm during the period of drying and setting.

MOLD PREVENTION

The risk of molding in bamboo culms can be reduced by drying culms such that the water content is reduced to less than 15 percent.

Stacking bamboo culms above the ground helps prevent molding and subsequent rotting. Storing bamboo in a cool and dry area also helps

reduce decay from molding. Coating the bamboo culms with borax or wood preservative chemicals inhibits the formation of molds.

PRESERVATION

Bamboo culms are susceptible to biological and physical deterioration especially when harvested young. Deterioration of the bamboo culm is mainly due to attack by powder-post beetles, termites, and decay caused by staining fungi. Methods to increase the durability or prolong the service life of bamboo culms are broadly classified into non-chemical and chemical methods.

NON-CHEMICAL METHODS

Many of the methods mentioned below are practiced traditionally and are suitable for small-scale industries or farmers with limited resources.

- **Curing:** After harvesting, the culms are left in the field for some time with branches and leaves intact. The transpiration of moisture through the leaves contributes to the reduction of starch in the culm.
- **Smoking:** The bamboo culms are cut into the desired length and stacked above a fire in an enclosed area. The smoke causes the culms to blacken and the heat destroys the starch in the parenchyma cells. Bamboo culms cured with smoke are known to last more than 15 years.
- **White Washing:** whole or split bamboo culms are painted with slaked lime. This prevents the entry of moisture into the culm, keeping away stain fungi and halting decay.
- **Construction Methods:** mounting bamboo poles over a concrete or stone foundation helps prolong their service life. Since the bamboo is not in contact with the ground, it is less susceptible to attack by fungi and termites.

- **Time of Harvesting:** harvesting mature (3 years old or older) culms during the dry season when their starch content is lowest makes them less vulnerable to attack by termites and fungi. If properly dried after harvesting, their useful life is significantly extended.
- **Plastering:** plastering bamboo culms or strips using cow dung mixed either with lime or mortar is effective in extending the durability of low-cost bamboo constructions.

CHEMICAL PRESERVATION METHODS


Using chemicals for preserving bamboo culms generally provides more effective protection than non-chemical methods. Chemical methods however imply greater costs and are only used when greater added-value and a higher quality product are required.

- **Fumigation:** involves the use of chemicals such as Methyl bromide for insect control.
- **Steeping or Sap Displacement:** green bamboo culms are allowed to stand vertically in a container of preservative solution till adequate chemical is picked up. At times, the culm may be freshly cut with branches and leaves on.
- **The Open-tank Treatment:** culms are cut to a desired length and are soaked in a solution of a water-soluble preservative for several days. The solution penetrates the culm by diffusion through the ends and partly through the sides.
- **Butt Treatment:** the bottom part of green bamboo or dried bamboo culm is immersed in a container of preservative, for example an old oil drum. The culms are left for about one week.
- **Old Engine Oil:** many farmers have been reported to use old engine oil particularly for green culms. The effectiveness of this method has not been widely reported and documented.

REFERENCES

1. Banik, R.L 1997. Domestication and Improvement of Bamboo. INBAR working paper No. 10. International Network for Bamboo and Rattan, New Delhi, India UNDP/ FAO forest and tree improvement project, Manila, the Philippines. 55.pp
2. Eggerling. W. J and Dale, I.R.1951. The Indigenous trees of the Uganda protectorate. Govern. Printers Entebbe, Uganda.
3. Hartmann, Handson T., Kester, Dale E. and Davies, Fred T. Jr. 1990 plant propagation principles and practices. Fifth edition. Prentice Hall, New Jersey. 647. pp.
4. Kigomo, B.N. 1988. Guidelines for the establishment and management of bamboo plantations in Kenya. Occasional paper No, 1: Kenya Forestry Research Institute. 31 pp.
5. Liese, W.1985. Bamboos-Biology, silvics, properties and utilization. GTZ. Germany, 132 pp.
6. LUSO CONSULT, 1997."Study on sustainable bamboo management" Final report on behalf of Deutsche Gesellschaft fur Technische Zusammenarbeit (GTZ) GmbH, Eschborn, Hamburg, 85 pp.
7. Marden, 1980. Bamboo the giant grass National Geographic. Vol.158.No.44, pp. 502-29.
8. McClure, F.A. 1996. The bamboo: A fresh perspective Harvard University press. Cambridge, Mass. Pp.347.
9. Mooney, H.F.1963.A glossary of Ethiopian plant names. Dublin Univer. Press Ltd.
10. Mooney, H.F. 1959. A report on the bamboo forests of Wellega Province with view to their possible utilization for pulp. Addis Ababa. p.8

11. Phillips. Sylvia. 1995. Flora of Ethiopia and Eritrea. Poaceae (Gramineae) National Herbarium, Addis Ababa University .p.5.
12. Wolde-Michael Kelecha, 1980. The bamboo potential of Ethiopia, FAWDA, Addis Ababa pp.14
13. Williamson, J.1974. Useful plants of Malawi. Pab.University Macawi, Indian.
14. Von Breitenbach, F.1963.The indigenous trees of Ethiopia, Addis Ababa, pp.195.



APPENDIX 1: DATA TO BE RECORDED BASED ON PHONOLOGICAL OBSERVATIONS OF BAMBOO

Phonological Events	Observations per Month											
	J	F	M	A	M	J	J	A	S	O	N	D
Flower bud formation												
Flowering												
Fruiting												
Matured fruits												
Fruit dropping												

APPENDIX 2: DATA TO BE GATHERED DURING VEGETATIVE PHASE OF HIGHLAND BAMBOO

Vegetative Phase	Observations per Month											
	J	F	M	A	M	J	J	A	S	O	N	D
Shoot Sprout												
Sheath Peel-off												
Clump Conditions After Reproductive Phase												
Rhizome Conditions After Reproductive Phase												
Interval Time Vegetative and Reproductive phase												

Symbol to be filled

SSH = Shoot Sprout

SSP = Sheath Peel –off

D= Die

L= Live